



Cambridge IGCSE™

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



ADDITIONAL MATHEMATICS

0606/22

Paper 2

February/March 2020

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Blank pages are indicated.

- 1 Find the values of x for which $12x^2 - 20x + 5 < (2x + 1)(x - 1)$. [4]



- 2 Variables x and y are such that, when $\lg y$ is plotted against x^3 , a straight line graph passing through the points $(6, 7)$ and $(10, 9)$ is obtained. Find y as a function of x . [4]



3 Find the exact solution of $3^{2x} - 3^{x+1} - 4 = 0$.


[4]



4 The position vectors of three points, A , B and C , relative to an origin O , are $\begin{pmatrix} -5 \\ -7 \end{pmatrix}$, $\begin{pmatrix} 10 \\ -4 \end{pmatrix}$ and $\begin{pmatrix} x \\ y \end{pmatrix}$ respectively. Given that $\overrightarrow{AC} = 4\overrightarrow{BC}$, find the unit vector in the direction of \overrightarrow{OC} .

[5]

- 5 (a) On the axes below, sketch the graph of $y = |5x - 7|$, showing the coordinates of the points where the graph meets the coordinate axes. [3]




- (b) Solve $5|5x - 7| - 1 = 14$. [3]

- 6 (a) A circle has a radius of 6 cm. A sector of this circle has a perimeter of $2(6 + 5\pi)$ cm. Find the area of this sector. [4]




(b)



The diagram shows the sector AOB of a circle with centre O and radius 7 cm. Angle $AOB = \frac{\pi}{4}$ radians. Find the perimeter of the shaded region. [3]

- 7 Find the coordinates of the points of intersection of the curves $x^2 = 5y - 1$ and $y = x^2 - 2x + 1$. [5]





The diagram shows the graph of $f(x) = a \cos bx + c$ for $0 \leq x \leq \frac{8\pi}{3}$ radians.

(a) Explain why f is a function. [1]

(b) Write down the range of f . [1]

(c) Find the value of each of the constants a , b and c . [4]

- 9 Variables x and y are such that $y = \frac{e^{3x} \sin x}{x^2}$. Use differentiation to find the approximate change in y as x increases from 0.5 to $0.5 + h$, where h is small. [6]

10 (a) $g(x) = 3 + \frac{1}{x}$ for $x \geq 1$.

7

(i) Find an expression for $g^{-1}(x)$.

[2]

(ii) Write down the range of g^{-1} .


[1]

(iii) Find the domain of g^{-1} .


[2]

(b) $h(x) = 2 \ln(3x - 1)$ for $x \geq \frac{2}{3}$.

The graph of $y = h(x)$ intersects the line $y = x$ at two distinct points. On the axes below, sketch the graph of $y = h(x)$ and hence sketch the graph of $y = h^{-1}(x)$. [4]



11



A container is a circular cylinder, open at one end, with a base radius of r cm and a height of h cm. The volume of the container is 1000 cm^3 . Given that r and h can vary and that the total outer surface area of the container has a minimum value, find this value. [8]

12 A particle P moves in a straight line such that, t seconds after passing through a fixed point O , its acceleration, $a \text{ ms}^{-2}$, is given by $a = -6t$. When $t = 0$, the velocity of P is 18 ms^{-1} .

(a) Find the time at which P comes to instantaneous rest. [3]

(b) Find the distance travelled by P in the 3rd second. [3]

13 (a) The sum of the first two terms of a geometric progression is 10 and the third term is 9.

7

(i) Find the possible values of the common ratio and the first term.

[5]

(ii) Find the sum to infinity of the convergent progression.

[1]

- (b) In an arithmetic progression, $u_1 = -10$ and $u_4 = 14$. Find $u_{100} + u_{101} + u_{102} + \dots + u_{200}$, the sum of the 100th to the 200th terms of the progression. [4]