



# Cambridge IGCSE™

CANDIDATE  
NAME

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--



**ADDITIONAL MATHEMATICS**

**0606/12**

Paper 1

**February/March 2021**

**2 hours**

You must answer on the question paper.

No additional materials are needed.

## INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.


## INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages.

- 1 Find the exact solutions of the equation  $3(\ln 5x)^2 + 2 \ln 5x - 1 = 0$ .

[4]

2  
7

The diagram shows the graph of  $y = a \sin bx + c$  where  $x$  is in radians and  $-2\pi \leq x \leq 2\pi$ , where  $a$ ,  $b$  and  $c$  are positive constants. Find the value of each of  $a$ ,  $b$  and  $c$ . [3]

3 The line  $AB$  is such that the points  $A$  and  $B$  have coordinates  $(-4, 6)$  and  $(2, 14)$  respectively.



(a) The point  $C$ , with coordinates  $(7, a)$  lies on the perpendicular bisector of  $AB$ . Find the value of  $a$ . [4]


(b) Given that the point  $D$  also lies on the perpendicular bisector of  $AB$ , find the coordinates of  $D$  such that the line  $AB$  bisects the line  $CD$ . [2]

- 4 (a) Show that  $2x^2 + 5x - 3$  can be written in the form  $a(x+b)^2 + c$ , where  $a$ ,  $b$  and  $c$  are constants. [3]



- (b) Hence write down the coordinates of the stationary point on the curve with equation  $y = 2x^2 + 5x - 3$ . [2]

- (c) On the axes below, sketch the graph of  $y = |2x^2 + 5x - 3|$ , stating the coordinates of the intercepts with the axes. [3]



- (d) Write down the value of  $k$  for which the equation  $|2x^2 + 5x - 3| = k$  has exactly 3 distinct solutions. [1]

5 In this question all lengths are in kilometres and time is in hours.



Boat  $A$  sails, with constant velocity, from a point  $O$  with position vector  $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$ . After 3 hours  $A$  is at the point with position vector  $\begin{pmatrix} -12 \\ 9 \end{pmatrix}$ .

(a) Find the position vector,  $\overrightarrow{OP}$ , of  $A$  at time  $t$ . [1]

At the same time as  $A$  sails from  $O$ , boat  $B$  sails from a point with position vector  $\begin{pmatrix} 12 \\ 6 \end{pmatrix}$ , with constant velocity  $\begin{pmatrix} -5 \\ 8 \end{pmatrix}$ .

(b) Find the position vector,  $\overrightarrow{OQ}$ , of  $B$  at time  $t$ . [1]

(c) Show that at time  $t$   $|\overrightarrow{PQ}|^2 = 26t^2 + 36t + 180$ . [3]

(d) Hence show that  $A$  and  $B$  do not collide. [2]

6 (a) A geometric progression has first term 10 and sum to infinity 6.



(i) Find the common ratio of this progression.

[2]

(ii) Hence find the sum of the first 7 terms, giving your answer correct to 2 decimal places. [2]

(b) The first three terms of an arithmetic progression are  $\log_x 3$ ,  $\log_x(3^2)$ ,  $\log_x(3^3)$ .

(i) Find the common difference of this progression. [1]

(ii) Find, in terms of  $n$  and  $\log_x 3$ , the sum to  $n$  terms of this progression. Simplify your answer. [2]


(iii) Given that the sum to  $n$  terms is  $3081 \log_x 3$ , find the value of  $n$ . [2]

(iv) Hence, given that the sum to  $n$  terms is also equal to 1027, find the value of  $x$ . [2]

## 7 DO NOT USE A CALCULATOR IN THIS QUESTION



In this question all lengths are in centimetres.



The diagram shows a trapezium  $ABCDE$  such that  $AB$  is parallel to  $EC$  and  $ABCD$  is a rectangle. It is given that  $BC = \sqrt{17} + 1$ ,  $ED = \sqrt{17} - 1$  and  $DC = \sqrt{17} + 4$ .

- (a) Find the perimeter of the trapezium, giving your answer in the form  $a + b\sqrt{17}$ , where  $a$  and  $b$  are integers. [3]

- (b) Find the area of the trapezium, giving your answer in the form  $c + d\sqrt{17}$ , where  $c$  and  $d$  are integers. [2]

(c) Find  $\tan AED$ , giving your answer in the form  $\frac{e+f\sqrt{17}}{8}$ , where  $e$  and  $f$  are integers. [2]

(d) Hence show that  $\sec^2 AED = \frac{81+9\sqrt{17}}{32}$ . [2]

8 (a) (i) Show that  $\sin x \tan x + \cos x = \sec x$ .

[3]



(ii) Hence solve the equation  $\sin \frac{\theta}{2} \tan \frac{\theta}{2} + \cos \frac{\theta}{2} = 4$  for  $0 \leq \theta \leq 4\pi$ , where  $\theta$  is in radians. [4]

(b) Solve the equation  $\cot(y + 38^\circ) = \sqrt{3}$  for  $0^\circ \leq y \leq 360^\circ$ .


[3]

9 The polynomial  $p(x) = 2x^3 - 3x^2 - x + 1$  has a factor  $2x - 1$ .



(a) Find  $p(x)$  in the form  $(2x - 1)q(x)$ , where  $q(x)$  is a quadratic factor.

[2]



The diagram shows the graph of  $y = \frac{1}{x}$  for  $x > 0$ , and the graph of  $y = -2x^2 + 3x + 1$ . The curves intersect at the points  $A$  and  $B$ .

(b) Using your answer to **part (a)**, find the exact  $x$ -coordinate of  $A$  and of  $B$ .

[4]

(c) Find the exact area of the shaded region.

[6]

10 A curve has equation  $y = \frac{(2x^2 + 10)^{\frac{3}{2}}}{x-1}$  for  $x > 1$ .

7

(a) Show that  $\frac{dy}{dx}$  can be written in the form  $\frac{(2x^2 + 10)^{\frac{1}{2}}}{(x-1)^2}(Ax^2 + Bx + C)$ , where  $A$ ,  $B$  and  $C$  are integers. [5]

(b) Show that, for  $x > 1$ , the curve has exactly one stationary point. Find the value of  $x$  at this stationary point. [4]