



Cambridge IGCSE™

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



ADDITIONAL MATHEMATICS

0606/13

Paper 1

October/November 2023

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS


- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages.

- 1 (a) On the axes, sketch the graphs of $y = 2x + 5$ and $y = |4x - 3|$, stating the intercepts with the coordinate axes. [3]




- (b) Solve the inequality $|4x - 3| < 2x + 5$. [3]

- 2 The perpendicular bisector of the line joining the points $\left(-3, \frac{2}{3}\right)$ and $\left(6, -\frac{7}{3}\right)$ passes through the point $(2, k)$. Find the value of k . [4]

3 On the axes, draw the graph of $y = 2 \sin \frac{x}{3} - 1$ for $-360^\circ \leq x \leq 360^\circ$.

[4]



4 The polynomial P is given by $P(x) = ax^3 + bx^2 + 3x + 2$, where a and b are integers. $P(x)$ has a factor of $2x + 1$. $P(x)$ has a remainder of -6 when divided by $x + 1$.



(a) Find the values of a and b .

[5]

(b) Show that the equation $P(x) = 0$ has only one real root.

[3]

- 5 (a) A 5-character password is to be formed from the following 10 characters.



Letters	A	B	C	X	Y	Z
Symbols	*	\$	#	&		

No character can be used more than once in any 5-character password.

- (i) Find the number of passwords that can be formed. [1]
- (ii) Find the number of passwords that can be formed if the password has to contain at least one symbol. [2]
- (iii) Find the number of passwords that can be formed if the password has to start with two letters and end with two symbols. [2]
- (b) A team of 8 people is to be chosen from 5 doctors, 4 teachers and 6 police officers.
Find how many possible teams have the same number of doctors as teachers. [5]

6 The polynomial $q(x)$ is given by $q(x) = -\frac{1}{3}(2x-1)(x+3)^2$.


7

(a) Find the x -coordinates of the stationary points on the curve $y = q(x)$.

[4]

(b) On the axes, sketch the graph of $y = q(x)$ stating the intercepts with the coordinate axes.

[3]



(c) Find the values of k such that $q(x) = k$ has exactly one solution.

[3]

7 Solve the equation $6x^{\frac{1}{3}} - 2x^{-\frac{1}{3}} - 1 = 0$. Give your answers in exact form.

[4]




- 8 The first three terms, in descending powers of x , in the expansion of $\left(2x^2 - \frac{1}{4x}\right)^n$ can be written in the form $256x^{16} + ax^{13} + bx^c$, where n, a, b and c are integers. Find the values of n, a, b and c . [6]

- 9 Given that $y = \frac{(5x+2)^{\frac{1}{3}}}{(x-1)^2}$, show that $\frac{dy}{dx}$ can be written in the form $\frac{-(Ax+B)}{3(5x+2)^{\frac{2}{3}}(x-1)^3}$, where A and B are integers. [5]

10 In this question, all lengths are in centimetres and all angles are in radians.

R




The diagram shows the sector, OAB , of a circle with centre O and radius 20. The perimeter of this sector is 65. The lines CA and CB are both tangents to the circle at the points A and B , so that the triangle ABC is isosceles, with $AC = CB$. The angle AOB is equal to θ .

Find the area of the shaded region.

[9]

11



In the triangle OAB , $\vec{OA} = \mathbf{a}$ and $\vec{OB} = \mathbf{b}$.

The straight line XYZ is such that:

- $\vec{OX} = \frac{4}{5}\mathbf{b}$
- $\vec{AY} = \frac{1}{3}\vec{AB}$
- $\vec{AZ} = \mu\mathbf{a}$, where μ is a constant
- $\vec{YZ} = \lambda\vec{XY}$, where λ is a constant.

(a) Show that $\vec{XY} = \frac{2}{3}\mathbf{a} - \frac{7}{15}\mathbf{b}$.

[3]

(b) Find \overrightarrow{YZ} in terms of λ , \mathbf{a} and \mathbf{b} . [1]

(c) Find \overrightarrow{YZ} in terms of μ , \mathbf{a} and \mathbf{b} . [2]

(d) Hence find the values of λ and μ , [3]

- 12 Solve the equation $3 \operatorname{cosec}^2\left(\frac{2x}{3} - \frac{\pi}{3}\right) = 4$, for $0 < x \leq 3\pi$. Give your answers in terms of π . [5]

7