

(a) (i) Translate triangle *T* by the vector  $\begin{pmatrix} -7 \\ 1 \end{pmatrix}$ . Label the image *K*. [2]

(ii) Describe fully the **single** transformation that maps triangle *K* onto triangle *T*.  
 ..... [1]

(b) Reflect triangle *T* in the line  $y = 4$ . [2]

(c) Rotate triangle *T* through  $90^\circ$  clockwise about  $(0, 0)$ . [2]

(d) (i) Enlarge triangle *T* by scale factor  $-\frac{1}{2}$ , centre  $(0, 0)$ . Label the image *P*. [2]

(ii) Describe fully the **single** transformation that maps triangle *P* onto triangle *T*.  
 ..... [2]

- 2 (a) Daisy records her 50 homework marks.  
 The table shows the results.



Homework mark	15	16	17	18	19	20
Frequency	1	3	19	11	10	6

- (i) Write down the range.  
 ..... [1]
- (ii) Write down the mode.  
 ..... [1]
- (iii) Find the median.  
 ..... [1]
- (iv) Calculate the mean.  
 ..... [3]

- (b) 21 33 20 25 21 34 22 21 20 30 18

The list shows Ed's scores in 11 tests.

- (i) Complete the stem-and-leaf diagram to show this information.

1	
2	
3	

Key: 2|5 represents 25

- (ii) Find the median.  
 ..... [1]
- (iii) Find the interquartile range.  
 ..... [2]

- 3 (a) The value of Priya's car decreases by 10% every year.  
The value today is \$7695.



- (i) Calculate the value of the car after one year.

\$ ..... [2]

- (ii) Calculate the value of the car one year ago.

\$ ..... [2]

- (b) Ali invests \$600 at a rate of 2% per year simple interest.

Calculate the value of Ali's investment at the end of 5 years.

\$ ..... [3]

- (c) Sara invests \$500 at a rate of  $r\%$  per year compound interest.  
At the end of 12 years, the value of Sara's investment is \$601.35, correct to the nearest cent.

Find the value of  $r$ .

$r =$  ..... [3]

(d) The mass of a radioactive substance decreases exponentially at a rate of 3% each day.



(i) Find the overall percentage decrease at the end of 10 days.

..... % [2]

(ii) Find the number of whole days it takes until the mass of this substance is one half of its original amount.

..... [3]

4 (a)


NOT TO  
SCALE

The diagram shows a sector of a circle that is made into a cone by joining  $OA$  to  $OB$ .  
The sector angle is  $x^\circ$  and the radius of the sector is 7.5 cm.  
The base radius of the cone is 1.5 cm.

Calculate the value of  $x$ .

$x = \dots\dots\dots$  [3]

(b)

NOT TO  
SCALE

The diagram shows a cylinder with radius 8 cm inside a sphere with radius 17 cm.  
Both ends of the cylinder touch the curved surface of the sphere.


(i) Show that the height of the cylinder is 30 cm.

[2]

(ii) Calculate the volume of the cylinder as a percentage of the volume of the sphere.

..... % [4]

(c)



NOT TO  
SCALE

The diagram shows a solid sphere with radius 6 cm inside a cube with side length 20 cm. The cube contains water to a depth of 15 cm. The sphere is removed.

Calculate the new depth of water in the cube.

..... cm [3]

- 5 (a) In a shop the cost of a fiction book is  $\$x$  and the cost of a reference book is  $\$(x+2)$ .  
 The cost of 11 fiction books is the same as the cost of 10 reference books.



Find the value of  $x$ .

$$x = \dots\dots\dots [2]$$

- (b) In another shop, the cost of a fiction book is  $\$y$  and the cost of a reference book is  $\$(y+2)$ .  
 Maria spends  $\$95$  on fiction books and  $\$147$  on reference books.  
 She buys a total of 12 books.

- (i) Show that  $6y^2 - 109y - 95 = 0$ .


[4]

- (ii) Factorise  $6y^2 - 109y - 95$ .

$$\dots\dots\dots [2]$$

- (iii) Find the value of  $y$ .


$$y = \dots\dots\dots [1]$$


6  
NOT TO  
SCALE

The diagram shows a right-angled triangle.

Find the value of  $w$ .

$w = \dots\dots\dots [7]$

7 (a) 



NOT TO SCALE

The diagram shows a right-angled triangle  $PQR$  on horizontal ground.  $X$  is vertically above  $R$  and the angle of elevation of  $X$  from  $P$  is  $21^\circ$ .  $XR = 2.8\text{ m}$  and  $RQ = 7.1\text{ m}$ .


(i) Calculate the angle of elevation of  $X$  from  $Q$ .

..... [2]

(ii) Calculate  $PQ$ .

..... m [3]

(b)




NOT TO SCALE

Calculate the acute angle  $KML$ .

Angle  $KML =$  ..... [3]

(c)



The area of triangle  $ABC$  is  $62.89 \text{ cm}^2$ .

(i) Show that angle  $BAC = 28.4^\circ$ , correct to 1 decimal place.

[2]


(ii) Calculate  $BC$ .

..... cm [3]


(iii)  $AB$  is extended to a point  $D$  such that angle  $BDC = 90^\circ$ .

Calculate  $BD$ .

..... cm [3]



Dice A



Dice B

The diagram shows two fair dice.  
 Dice A is numbered 1, 2, 2, 2, 3, 6.  
 Dice B is numbered 2, 3, 3, 4, 4, 4.

(a) (i) Dice A is rolled once.

Write down the probability that it lands on the number 6.

..... [1]

(ii) Dice A is rolled 150 times.

Find the number of times it is expected to land on the number 6.

..... [1]

(b) Dice A and Dice B are each rolled once.

(i) Find the probability that the two numbers they land on have a total of 6.

..... [3]

(ii) Find the probability that when the two numbers they land on have a total of 6, both numbers are 3.

..... [2]


(c) Dice B is rolled  $n$  times.

The probability that on the  $n$ th roll it first lands on a number 3 is  $\frac{32}{729}$ .

Find the value of  $n$ .

$n = \dots\dots\dots$  [2]

9  

The diagram shows a sketch of the graph of  $y = 4x^3 - x^4$ .  
 The graph crosses the  $x$ -axis at the origin  $O$  and at the point  $A$ .  
 The point  $B$  is a maximum point.

(a) Differentiate  $4x^3 - x^4$ .

..... [2]

(b) Find the coordinates of  $B$ .

(....., .....) [3]

(c) Find the gradient of the graph at the point  $A$ .

..... [3]

11       $f(x) = 1 - 3x$        $g(x) = (x - 1)^2$        $h(x) = \frac{3}{x}, x \neq 0$



(a) Find  $g(3)$ .

..... [1]

(b) Find  $f(x-2)$ , giving your answer in its simplest form.

..... [2]

(c) Find  $f^{-1}(x)$ .

$f^{-1}(x) =$  ..... [2]

(d)  $gf(x) - g(x)f(x) = 3x^3 + ax^2 + bx + c$

Find the value of each of  $a$ ,  $b$  and  $c$ .

$a =$  .....

$b =$  .....

$c =$  ..... [5]

(e) Find  $h(x) - f(x)$ , giving your answer as a single fraction in its simplest form.


..... [3]

(f)  $h(x^n) = 3x^7$

Find the value of  $n$ .

$n =$  ..... [1]

12



NOT TO SCALE

$O$  is the origin  $(0, 0)$ ,  $A$  is the point  $(8, 1)$  and  $B$  is the point  $(2, 5)$ .

(a) Write as column vectors.

(i)  $\vec{OB}$

$$\vec{OB} = \begin{pmatrix} \phantom{0} \\ \phantom{0} \end{pmatrix} \quad [1]$$

(ii)  $\vec{AB}$

$$\vec{AB} = \begin{pmatrix} \phantom{0} \\ \phantom{0} \end{pmatrix} \quad [1]$$

- (b) Find the equation of the line  $AB$ .  
Give your answer in the form  $y = mx + c$ .

$$y = \dots\dots\dots [3]$$

- (c) Find the equation of the perpendicular bisector of  $AB$ .  
Give your answer in the form  $y = mx + c$ .

$$y = \dots\dots\dots [4]$$

- (d) The line  $AB$  meets the  $y$ -axis at  $P$ .  
The perpendicular bisector of  $AB$  meets the  $y$ -axis at  $Q$ .

Find the length of  $PQ$ .

$$\dots\dots\dots [2]$$